

## NOTA

---

 RODENT CONSUMPTION BY  
*PHILODRYAS PSAMMOPHIDEA*  
 (SERPENTES: COLUBRIDAE), FROM  
 THE INTER-ANDEAN DRY VALLEYS  
 OF CENTRAL BOLIVIA
 

---

OLIVER QUINTEROS-MUÑOZ

DIEGO A. PEÑARANDA

FREDDY NAVARRO

Centro de Biodiversidad y Genética, Universidad Mayor de San Simón, Casilla 538. Cochabamba, Bolivia.

ohlisin@yahoo.es

The genus *Philodryas* (Wagler, 1830) is composed by opisthoglyphous snakes with terrestrial and diurnal habits, some may also be arboreal. They are medium to large hunters with thin bodies (Thomas, 1976; Ceí, 1993, López and Girauo, 2008). Widely distributed in South America, this genus includes 18 species (Zaher *et al.*, 2008), of which eight have been recorded in Bolivia (Aguayo, 2009).

These snakes feed on a wide variety of small vertebrates, such as lizards, anurans, snakes, birds and small mammals even bats (Thomas, 1976; Funk *et al.*, 2003; Lopez, 2003; Otto and Miller, 2004; Franca and Araujo, 2007). However, the feeding habits of only two species, *P. patagoniensis* and *P. olfersii*, have been described in detail (Hartmann and Marques, 2005; Lopez and Girauo, 2008; Leite *et al.*, 2009). In this sense, *P. psammophidea* is regarded a herpetophagous snake (Leynaud *et al.*, 2006; Aguayo *et al.*, 2007; Franca and Araujo, 2007).

In May 18, 2009 we found an adult female of *Philodryas psammophidea* (930 mm SVL), at a side of a crop field in the Tabacal valley (18°23'7.42" S – 64°38'7.88" W, 2015 m), Narciso Campero province southern Cochabamba, Bolivia. Ecologica-

lly, this valley belongs to the Inter-Andean Dry Forests of Bolivia (Ibisch and Merida, 2003). In the stomach of the snake probably killed by a settler, there was an adult female of *Graomys domorum* (Phyllotini; Sigmodontidae), a native rodent species widely distributed in the region. This rodent was ingested head-first recently, because there were no signs of decomposition. Both specimens were deposited in the vertebrate collection of the Centro de Biodiversidad y Genética (*P. psammophidea* CBG-700.; *G. domorum* CBG-405).

Despite *Philodryas* members are considered generalists predators, amphibians and reptiles constitute the most consumed preys, while birds and small mammals occur less frequently and have been reported in the diet of only five species of the genus (*P. aestiva*, *P. baroni*, *P. nattereri*, *P. olfersii*, *P. patagoniensis*) (Ceí, 1993; Franca and Araujo, 2007). Although *P. psammophidea* is fed with rodents in captivity (Scrocchi *et al.*, 2006), this is the first formal evidence of small mammals consumption by this snake in wild in Bolivia, and suggests a generalist diet for *P. psammophidea* as observed in other snakes of the genus.

On the other hand, it is known that the feeding habits of the two best-studied species of the genus, *P. olfersii* and *P. patagoniensis*, vary geographically in the dominance of prey types they consume and it has been proposed that such changes may be related to the increased prey availability or detectability, the snakes activity patterns or the phylogenetic relationships among them, instead of a resource preference pattern, as the most consumed prey do not constitute limited resources in the environment (Hartmann, 2001, Hartmann and Marques, 2005, Lopez and Girauo, 2008, Leite *et al.*, 2009).

Therefore, considering that rodents are apparently more abundant elements

compared with amphibians and lizards in the inter Andean dry valleys southern Cochabamba (Cahill *et al.*, 2010), *P. psammophidea* could be an important controller of rodents, mainly in those areas close to villages and crop fields where the higher abundances have been observed (D. A. Peñaranda, unpub. data).

#### ACKNOWLEDGEMENTS

This work is a result of the project Technical Justification for the Establishment of the Pasorapa Integrated Managed Natural Area, funded by American Bird Conservancy and executed by the Center for Biodiversity and Genetics. Authors want to thank to Jennifer R.A. Cahill for her support and encouragement as the general coordinator of the project. To G. Scrocchi whose comments and suggestions helped to improve previous versions of this manuscript. To E. Rocha who found and collected the specimen, O. Osco whose experience with Andean rodents was important, finally to Adam W. Ferguson for providing literature of *Philodryas* snakes.

#### LITERATURE CITED

- AGUAYO, R. 2009. Reptiles de Bolivia. Bolivia Ecológica. Centro de Ecología y Difusión Simon I. Patiño 53: 32 pp.
- AGUAYO, R.; G. REY & O. RUIZ. 2007. Anfibios y Reptiles. 103-158, *En*: AGUIRRE, L. F.; R. AGUAYO; O. RUIZ & F. NAVARRO. (eds.), *Guía de los Mamíferos, Anfibios y Reptiles del Parque Nacional Tunari*. Centro de Biodiversidad y Genética, UMSS, Cochabamba, Bolivia.
- CAHILL, J. R. A.; A. ZEBALLOS; E. ROCHA; N. DE LA BARRA; J. M. LAZCANO; A. SELAYA & M. FERNÁNDEZ. 2010. Justificación para la creación del Área Natural de Manejo Integrado Municipal Pasorapa (ANMI Municipal Pasorapa). Centro de Biodiversidad y Genética, Universidad Mayor de San Simón, Cochabamba-Bolivia.
- CEI, J. M. 1993. Reptiles del noroeste, nordeste y este de la Argentina. *Museo Regionale Sci. Naturale Torino, Monografie* 14: 1-949.
- FRANCA, F. G. R. & A. F. B. ARAUJO. 2007. Are there co-occurrence patterns that structure snake communities in Central Brazil? *Brazilian Journal of Biology*, 67 (1): 33-40.
- FUNK, P. G.; M. DI-BERNARDO & M. BORGES-MARTINS. 2003. *Philodryas patagoniensis* (Papa-pinto), Cannibalism. *Herpetological Review* 34 (2): 154.
- GONZÁLEZ, L. & S. REICHLER. 2003. Anexo 3. Lista de reptiles presentes en Bolivia. Pp. 586-589, *In*: IBISCH, P. L. & G. MÉRIDA (eds.) Biodiversidad: la riqueza de Bolivia: estado de conocimiento y conservación. Editorial FAN, Santa Cruz de la Sierra.
- HARTMANN, P. A & O. A. V. MARQUES, 2005. Diet and habitat use of two sympatric species of *Philodryas* (Colubridae), in south Brazil. *Amphibia-Reptilia*. 26: 25-31.
- HARTMANN, P. A. 2001. Hábito alimentar e utilização do ambiente em duas espécies simpátricas de *Philodryas* (Serpentes, Colubridae), no sul do Brasil. Unpublished Master Thesis, Universidade Estadual Paulista, Rio Claro.
- IBISCH, P. L. & G. MERIDA. 2003. Biodiversidad: La riqueza de Bolivia. Estado de conocimiento y conservación. Ministerio de Desarrollo sostenible. Editorial FAN, Santa Cruz de La Sierra, Bolivia. 638 pp.
- LEITE, P. T.; I. L. KAEFER & S. Z. CENCHIN. 2009. Diet of *Philodryas olfersii* (Serpentes, Colubridae) during hydroelectric dam flooding in

- southern Brazil. *North-Western Journal of Zoology*. 5 (1): 53-60.
- LEYNAUD, G. C.; N. PELEGRIN & J. N. LESCANO. 2006. Anfibios y Reptiles: 219-235. In: BUCHER, E.H. (ed), Bañados del río Dulce y Laguna Mar Chiquita, Córdoba, Argentina. Academia Nacional de Ciencias (Córdoba, Argentina).
- LOPEZ, M. S. & A. R. GIRAUDO. 2008. Ecology of the Snake *Philodryas patagoniensis* (Serpentes, Colubridae) from Northeast Argentina. *Journal of Herpetology*, 42 (3): 474-480.
- LOPEZ, M. S. 2003. *Philodryas patagoniensis*. Diet. *Herpetological Review* 34 (1): 71-72.
- OTTO, C. R. & G. J. MILLER. 2004. *Philodryas viridissimus* (Green Palm Snake), Feeding attempt. *Herpetological Review* 35 (3): 227.
- SCROCCHI, G.; J. C. MORETA & S. KRETZCHMAR. 2006. Serpientes del Noroeste Argentino. Fundación Migue Lillo. Tucumán, Argentina, 174 pp.
- THOMAS, R. A. 1976. A revision of the South American snake genus *Philodryas* Wagler, 1830. Unpublished Ph.D. Thesis Texas A&M University. 338 pp.
- ZAHER, H.; G. SCROCCHI & R. MASIERO. 2008. Rediscovery and redescription of the type of *Philodryas laticeps* Werner, 1900 and the taxonomic status of *P. oligolepis* Gomes, 1921 (Serpentes, Colubridae). *Zootaxa*, 1940: 25-40.