

# First record of *Eustrongylides* sp. (Nematoda: Dioctophymatidae) larvae parasitizing snakes in the Neotropical region

Julia Veiga Pereira<sup>1</sup>, Roger Jean Oliveira<sup>2</sup>, Carolina Silveira Mascarenhas<sup>2</sup>, Gertrud Müller<sup>2</sup>

<sup>1</sup>Laboratório de Herpetologia, Instituto de Biociências, Universidade Federal do Rio Grande do Sul, Av. Bento Gonçalves, 9500, Agronomia, CEP 91501-970, Porto Alegre, Rio Grande do Sul, Brazil.

<sup>2</sup>Laboratório de Parasitologia de Animais Silvestres, Instituto de Biologia, Universidade Federal de Pelotas, Campus Universitário, CEP 96010-900, Pelotas, Rio Grande do Sul, Brazil.

Recibida: 25 Mayo 2020

Revisada: 08 Julio 2020

Aceptada: 15 Septiembre 2020

Editor Asociado: C. Borteiro

doi: 10.31017/CdH.2020.(2020-042)

## ABSTRACT

*Eustrongylides* uses piscivorous birds as definitive hosts, oligoquets as intermediate hosts and fish as second intermediate hosts. However, amphibians and reptiles can act as secondary or paratenic hosts. The following work presents the first record of *Eustrongylides* larvae in snakes in the Neotropics. The coelomic cavity of 101 snakes belonging to *Helicops infrataeniatus*, *Philodryas aestiva*, *Philodryas olfersii*, *Philodryas patagoniensis*, *Erythrolamprus poecilogyrus*, *Erythrolamprus jaegeri*, *Atractus reticulatus*, *Dipsas ventrimaculatus*, *Thamnodynastes strigatus*, *Phalotris lemmiscatus* and *Bothrops alternatus* from southern Brazil were examined. Six *H. infrataeniatus* were parasitized with larvae of *Eustrongylides* sp. The encounter of these larvae only in this specie can be related to the aquatic habit of the snake, which allows ecological interactions with the definitive and intermediate hosts of the *Eustrongylides* cycle. The occurrence of these parasites in a Neotropical snake is registered for the first time, contributing to the knowledge of the helminth diversity associated to Brazilian snakes.

Key Words: *Helicops infrataeniatus*; Helminths; Parasitism; Reptiles; Serpentes.

*Eustrongylides* (Jägerskiöld, 1909) (Nematoda: Dioctophymatidae) was proposed for Dioctophymatidae species that cause tumors on the wall of the proventriculus, mainly from piscivorous birds (Measures, 1988). Some species were reported in Gaviiformes, Podicipediformes, Pelecaniformes, Anseriformes, Charadriiformes, Ciconiiformes and Passeriformes (Measures, 1988). *Eustrongylides* consists of three species: *Eustrongylides tubifex* Jägerskiöld, 1909, *Eustrongylides ignotus* (Jägerskiöld, 1909) and *Eustrongylides excisus* (Jägerskiöld, 1909) (Melo *et al.*, 2016).

Regarding the geographical distribution of these parasites, *Eustrongylides ignotus* has the widest distribution, occurring in the Nearctic, Neotropical and Australian regions. *Eustrongylides excisus* occurs in Europe and *E. tubifex* is present only in the

Holarctic and Neotropical regions (Measures, 1988).

The life cycle of these parasites is not well known (Xiong *et al.*, 2009), however, some studies indicate that these helminths use piscivorous birds as definitive hosts, oligoquets as intermediate hosts and usually fish as intermediate second hosts, with amphibians and reptiles as intermediate or paratenic hosts (Measure, 1988; Friend and Franson, 1999; Melo *et al.*, 2016). *Eustrongylides* species may occasionally parasitize humans, thus demonstrating zoonotic potential (Xiong *et al.*, 2009; Eiras *et al.*, 2016).

Little is known about the parasitism of *Eustrongylides* species in wild snake populations, but larvae of these parasites have been reported in some Colubridae from North America, Europe and Asia,

such as: *Nerodia sipedon* (Linnaeus, 1758), *Natrix natrix* (Linnaeus, 1751) and *N. tessellata* (Laurenti, 1768) (Burse, 1986; Biserkov, 1995; Mihalca, 2007; Yildirimhan *et al.*, 2007). In acrochordid snakes from Australia there are also reports of the occurrence of these parasites (Jones, 1978). In captivity, larvae of *Eustrongylides* have been observed in some colubrid snakes such as *Coluber constrictor* (Linnaeus, 1758), *Masticophis flagellum* (Shaw, 1802), *Nerodia sipedon*, *Pituophis melanoleucus* (Daudin, 1803), *Thamnophis sirtalis* (Linnaeus, 1758), as well as in the viperid snakes *Bothrops atrox* (Linnaeus, 1758) and *Agkistrodon contortrix* (Linnaeus, 1766) in North America (Winsor, 1948 in Bursey, 1986). In Brazil, while these nematodes have been reported in some hosts, such as birds (Vicente *et al.*, 1995; Bernardon *et al.*, 2017), amphibians (Melo *et al.*, 2016) and fish (Isaac *et al.*, 2004; Meneguetti *et al.*, 2013; Carvalho *et al.*, 2017), we know of no reports of their occurrence in snakes. The following work presents the first record of *Eustrongylides* larvae in wild snakes from the Neotropical region.

One hundred and one specimens belonging to 11 species of snakes were examined: *Helicops infrataeniatus* Jan, 1865 (n=65), *Philodryas aestiva* (Duméril, Bibron & Duméril, 1854) (n=1), *P. olfersii* (Lichtenstein, 1823) (n=4), *P. patagoniensis* (Girard, 1858) (n=6), *Erythrolamprus poecilogyrus* (Wied-Neuwied, 1825) (n=10), *E. jaegeri* (Günther, 1858) (n=2), *Atractus reticulatus* (Boulenger, 1885) (n=1), *Dipsas ventrimaculatus* (Boulenger, 1885) (n=3), *Thamnodynastes strigatus* (Günther, 1858) (n=2), *Phalotris lemniscatus* (Duméril, Bibron & Duméril, 1854) (n=1) (Dipsadidae) and *Bothrops alternatus* (Duméril, Bibron & Duméril, 1854) (n=6) (Viperidae).

The snakes came from the municipalities of Capão do Leão (31° 45' 48" S - 52° 29' 02" W), Pelotas (31° 46' 19" S - 52° 20' 33" W), Rio Grande (32° 02' 06" S - 52° 05' 55" W), Encruzilhada do Sul (30° 32' 38" S - 52° 31' 19" W) and Dom Pedrito (30° 58' 58" S - 54° 40' 23" W), Rio Grande do Sul. Fifty of them were collected dead on roads from March 2017 to June 2019. The collection of specimens were licensed by Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio No. 38913). Four snakes were donated by the Núcleo de Reabilitação da Fauna Silvestre e Centro de Triagem de Animais Silvestres of the Federal University of Pelotas (NURFS-CETAS/UFPel), where they died after a rehabilitation attempt. The other hosts were donated by the Ver-

tebrate Laboratory of the Federal University of Rio Grande (n= 36) and Vertebrate Zoology Laboratory of the Federal University of Pelotas (n= 11), where the specimens were fixed in formalin and conserved in 70°GL alcohol.


The hosts were necropsied for analysis of the coelomic cavity. The larvae were fixed in AFA, conserved in 70°GL alcohol and clarified with Amann lactophenol for morphological identification according to Measures (1988). The parasitological indices were calculated according to Bush *et al.* (1997). The specimens were deposited in the helminth collection of the Laboratory of Wild Animal Parasitology (CHLAPASIL n° 831-836). The photomicrographs were done in an Olympus® BX 41 microscope with attached camera system and the images were prepared in Adobe Photoshop® CS5.

Among the total hosts examined, six (5.9%) were parasitized by fourth stage larvae of *Eustrongylides* sp. (Fig. 1), which were found in the coelomic cavity of *Helicops infrataeniatus* (Fig. 2), with a prevalence of 9.2% and mean intensity of infection of 1.83 helminths/host. Female larvae were larger than male larvae, which measured 75-115 mm and 65-80 mm length, respectively.

*Helicops infrataeniatus* is a non-venomous aquatic snake that can reach about one meter in length. It is distributed in southern Brazil, Paraguay, Uruguay and Argentina (Quintela and Loebmann, 2009; Abegg and Entiauspe-Neto, 2012). The diet of the species includes mostly fish and amphibians (Aguiar and Di-Bernardo, 2004), which were possibly the source of infestation of the snakes in this study.

The transmission of *Eustrongylides* species occurs through the trophic chain involving aquatic organisms such as oligochaetes (primary intermediate hosts), fish, anurans, snakes (secondary and/or paratenic intermediate hosts) and birds (definitive hosts). Bird infection occurs by ingestion of the infective fourth-stage larva present in the secondary and/or paratenic intermediate hosts (Measure, 1988; Moravec, 1994; Friend and Franson, 1999).

Larvae are often found parasitizing several fish species in Brazil, such as *Hoplias malabaricus* (Bloch, 1794) (Characiformes), *Cichla piquiti* Kullander & Ferreira, 2006 and *Plagioscion squamosissimus* (Heckel, 1840) (Perciformes) (Isaac *et al.*, 2004; Martins *et al.*, 2009; Meneguetti *et al.*, 2013; Carvalho *et al.*, 2017). Amphibians were also reported as hosts of *Eustrongylides* larvae in northern Brazil (Melo *et*




**Figure 1.** Fourth stage larvae from *Eustrongylides* sp. (Enoplida: Dioctophymatidae) found in the coelomic cavity of *Helicops infrataeniatus* Jan, 1865 (Serpentes: Dipsadidae) in the extreme south of Brazil. A - Anterior extremity of female larva (Arrows indicate the cephalic papillae; oe: esophagus; bar - 100  $\mu$ m). B - Posterior extremity of the female larva (Arrows indicate the primordial vulvar; bar - 100 $\mu$ m). C - Arrows indicate cuticle layers (oe: esophagus; bar - 12,5  $\mu$ m). D - Posterior end of the male larva (bar - 60  $\mu$ m).

al., 2016). There are no previous records of larvae of this nematode associated with snakes in Brazil; however, there are records in North America (Burse, 1986), Europe (Mihalca *et al.*, 2007; Biserkov, 1995; Yildirimhan *et al.*, 2007) and Australia (Lichtenfels and Lavies, 1976; Jones, 1978).

Snakes can include in their diet of large variety of vertebrates (Voris and Murphy, 2002; Abegg *et al.*, 2005; Costa *et al.*, 2009), enabling the transmission of parasites through complex trophic networks (Marcogliese, 2004). Herons, such as *Syrigma sibilatrix* (Temminck, 1824) (Ardeidae) feed on *Helicops infrataeniatus* as observed by Franz *et al.* (2007) in Campo Belo do Sul in the state of Santa Catarina, Brazil. In the region of our present study, *Eustron-*

*gylides ignotus* was reported in herons *Ardea cocoi* Linnaeus, 1766 and *Nycticorax nycticorax* (Linnaeus, 1758) (Ardeidae) (Bernardon *et al.*, 2017), and larvae of this nematode were found in siluriform fish, *Callichthys callichthys* (Linnaeus, 1758), as well as in *Hoplias malabaricus* (Carolina Silveira Mascarenhas, com. pess.). The record of fourth stage larvae of *Eustrongylides* sp. in *H. infrataeniatus* suggests that the species may act as a trophic bridge in the parasite cycle, since the snake has aquatic habits, feeds on fish and serves as a food resource for birds such as herons. These records highlight the importance of the region for maintenance of the *Eustrongylides* species life cycle and warns about the presence of a nematode with zoonotic potential (Eiras *et al.* 2016).



**Figure 2.** *Eustrongylides* sp. fourth stage larva (Enoplida: Dioctophymatidae) found in the coelomic cavity of *Helicops infrataeniatus* Jan, 1865 (Serpentes: Dipsadidae) in the extreme south of Brazil (Bar – 12 mm).

It is important to point out that the other snakes used in this study present predominantly terrestrial habits, such as *Bothrops alternatus*, *Thamnodynastes strigatus*, *Philodryas patagoniensis*, or semi-aquatic habit as *Erythrolamprus jaegeri* and *E. poecilogyrus* (Achaval and Olmos, 2003), which may hinder or even make completely impossible the transmission of this nematode. However, future studies should be conducted to complement the results presented herein to help in the understanding of parasitic relationships involving *Eustrongylides* species and snakes in the Neotropical region.

The occurrence of *Eustrongylides* larvae in *Helicops infrataeniatus* registered for the first time, is the second finding of helminths parasitizing this species in Brazil, since there is a previous record of parasitism by *Opisthogonimus serpentis* Artigas, Ruiz & Leão 1943 (Digenea: Opistogonimidae) (Fernandes and Kohn, 2014).

### Acknowledgments

Special thanks to José Dornelles, Daniel Loebmann, Rafael Dias, Marco Coimbra, Juliana Wolter and Ricardo da Silva for their assistance. This work was supported by CAPES (Coordenação de Aperfeiçoamento do Pessoal de Nível Superior, process number 32/2010) and CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico) through scholarship to JVP.

### Literature cited

- Abegg, A.D.; Entiauspe-Neto, O.M. 2012. Serpentes do Rio Grande do Sul. Tapera: Livraria & Editora Werlang.
- Abegg, A.D.; Entiauspe-Neto, O.M.; Costa, H.C. & Santos, P.S. 2015. *Erythrolamprus poecilogyrus* sspp. (Serpentes: Dipsadidae): predação. *Herpetologia Brasileira* 4: 60-63.
- Achaval, F. & Olmos, A. 2003. Anfíbios y Reptiles del Uruguay. Graphis, Montevideo.
- Aguiar, L.F.S. & Di-Bernardo, M. 2004. Diet and feeding behavior of *Helicops infrataeniatus* (Serpentes: Colubridae).

- Xenodontinae) in Southern Brazil. *Studies on Neotropical Fauna and Environment* 39: 7-14.
- Bernardon, F.F.; Valente, A.L.S. & Müller, G. 2017. Helminth assemblage of aquatic birds (Pelecaniformes: Ardeidae) of southern Rio Grande and a checklist of helminths of herons of Brazil. *Neotropical Helminthology* 11: 357-375.
- Biserkov, V.Y. 1995. New records of nematodes and acanthocephalans from snakes in Bulgaria. *Comptes Rendus de l'Académie Bulgare des Sciences* 48: 87-89.
- Burse, C.R. 1986. Histological aspects of natural eustrongyloid infections of the northern water snake, *Nerodia sipedon*. *Journal of Wildlife Diseases* 22: 527-532.
- Bush, A.O.; Lafferty, K.; Lotz, J. & Shostak, A. 1997. Parasitology meets ecology on its own terms: Margolis et al. revisited. *The Journal of Parasitology* 83: 575-583.
- Carvalho, A.R.; Martins, R.T.; Bellei, P.M. & Lima, S.S. 2017. Aspectos ecológicos da helmintofauna de *Hoplias malabaricus* (Bloch, 1794) (Characiformes, Erythrinidae) da Represa Dr. João Penido (Juiz de Fora-MG, Brasil). *Revista Brasileira de Zootecnia* 18: 7-20.
- Costa, H.C.; Assis, C.L.; Werneck, H.A.; Nunes, A.V. & Feio, R.N. 2009. Ataque de um falcão acauã (*Herpetotheres cachimans*) sobre a serpente boipeva (*Xenodon merremii*) na Mata Atlântica de Minas Gerais, sudeste do Brasil. *Revista Brasileira de Zootecnia* 11: 171-173.
- Eiras, J.C.; Velloso, A.L. & Pereira-Jr, J. 2016. Parasitos de peixes marinhos da América do Sul. Editora da FURG. Rio Grande.
- Fernandes, B.M.M. & Kohn, A. 2014. South American trematodes parasites of amphibians and reptiles. Oficina de Livros. Rio de Janeiro.
- Franz, I.; Ghizoni-Jr, I.R., Albuquerque, J.L.B.; Barcellos, A.; Hassdenteufel, C.B.; Arend, F.L. & Martins-Ferreira, C. 2007. Predação da cobra d'água *Helicops infrataeniatus* (Serpentes, Colubridae) pela maria-faceira *Syrigma sibilatrix* (Aves, Ardeidae) no sul do Brasil. *Biotemas* 20: 135-137.
- Friend, M. & Franson, J.C. 1999. Field manual of wildlife diseases. General field procedures and diseases of birds. Geological Survey, Biological Resources Division. Washington DC.
- Isaac, A.; Guidelli, G.M.; França, J.G. & Pavanelli, G.C. 2004. Composição e estrutura das infracomunidades endoparasitárias de *Gymnotus* spp. (Pisces: Gymnotidae) do rio Baía, Mato Grosso do Sul, Brasil. *Acta Scientiarum. Biological Sciences* 26: 453-462.
- Jones, H.I. 1978. Gastrointestinal nematodes from aquatic Australian snakes. *Memoirs of the Queensland Museum* 18: 243-254.
- Lichtenfels, J.R. & Lavies, B. 1976. Mortality in red-sided garter snakes, *Thamnophis sirtalis parietalis*, due to larval nematode, *Eustrongylides* sp. *Laboratory Animal Science* 26: 465-467.
- Marcogliese, D.J. 2004. Parasites: small players with crucial roles in the ecological theater. *EcoHealth* 1: 151-164.
- Martins, M.L.; Santos, R.D.S.; Marengoni, N.G.; Takahashi, H.K. & Onaka, E.M. 2009. Seasonality of *Eustrongylides* sp. (Nematoda: Dioctophymatidae) larvae in fishes from Paraná River, South-western Brazil. *Boletim do Instituto de Pesca* 35: 29-37.
- Measures, L.N. 1988. The development of *Eustrongylides tubifex* (Nematoda: Dioctophymatoidea) in oligochaetes. *Journal of Parasitology* 74: 294-304.
- Melo, F.T.V.; Melo, C.S.B.; Nascimento, L.C.S.; Giese, E.G.; Furtado, A.P. & Santos, J.N. 2016. Morphological characterization of *Eustrongylides* sp. larvae (Nematoda, Dioctophymatoidea) parasite of *Rhinella marina* (Amphibia: Bufonidae) from Eastern Amazonia. *Revista Brasileira de Parasitologia Veterinária* 25: 235-239.
- Meneguetti, D.U.O.; Laray, M.P.O. & Camargo, L.M.A. 2013. Primeiro relato de larvas de *Eustrongylides* sp. (Nematoda: Dioctophymatidae) em *Hoplias malabaricus* (Characiformes: Erythrinidae) no Estado de Rondônia, Amazônia Ocidental, Brasil. *Revista Pan-Amazônica de Saúde* 4: 55-58.
- Mihalca, A.D.; Fictum, P.; Škorič, M.; Sloboda, M.; Kärverno, S.; Ghira, I., Carlsson, M. & Modrý, D. 2007. Severe granulomatous lesions in several organs from *Eustrongylides* larvae in a free-ranging dice snake, *Natrix tessellata*. *Veterinary pathology* 44: 103-105.
- Moravec, F. 1994. Parasitic nematodes of freshwater fishes of Europe. Kluwer Academic Publishers. Netherlands.
- Quintela, F.M. & Loebmann, D. 2009. Os répteis da região costeira do extremo sul do Brasil: guia ilustrado. USEB. Pelotas.
- Vicente, J.J.; Pinto, R.M.; Noronha, D. & Gonçalves, L. 1995. Nematode parasites of Brazilian Ciconiiformes birds: a general survey with new records for the species. *Memórias do Instituto Oswaldo Cruz* 90: 389-393.
- Voris, H.K. & Murphy, J.C. 2002. The prey and predators of homalopsine snakes. *Journal of Natural History*, 36: 1621-1632.
- Xiong, F.; Wang, G.T.; Wu, S.G. & Nie, P. 2009. Development of *Eustrongylides ignotus* (Nematoda: Dioctophmida) in Domestic Ducks (*Anas platyrhynchos domestica* (L.)). *The Journal of Parasitology* 95: 1035-1039.
- Yildirimhan, H.S.; Bursey, C.R. & Goldberg, S.R. 2007. Helminth parasites of the grass snake, *Natrix natrix*, and the dice snake, *Natrix tessellata* (Serpentes: Colubridae), from Turkey. *Comparative Parasitology* 74: 343-354.

