

## Predation on *Abronia graminea* (Squamata: Anguidae) and *Sceloporus bicanthalis* (Squamata: Phrynosomatidae) by *Lanius ludovicianus* (Laniidae, Aves) in Veracruz, Mexico

Víctor Vásquez-Cruz, Arleth Reynoso-Martínez

<sup>1</sup>PIMVS Herpetario Palancoatl, Avenida 19 número 5525, Colonia Nueva Esperanza, C.P. 94540, Córdoba, Veracruz, México

Recibida: 25 Marzo 2020  
Revisada: 09 Julio 2020  
Aceptada: 14 Septiembre 2020  
Editor Asociado: J. Goldberg

doi: 10.31017/CdH.2020.(2020-046)

### ABSTRACT

Here we document the first case of predation on the lizards *Abronia graminea* and *Sceloporus bicanthalis*, presumably by the Loggerhead Shrike, *Lanius ludovicianus*, in west-central Veracruz, Mexico

Key Words: Diet, impalement behavior, Loggerhead Shrike, predator-prey interactions.

Predation can be an important factor in the evolution of species' life history traits (e. g. Steen *et al.*, 2014). However, is rarely observed in the wild and few studies have confirmed the identity of predators and preys either by direct observation of predation events, or by traces of the interaction, such as traces of prey in feces (e.g. Vásquez-Cruz and Reynoso-Martínez, 2020) or stomach dissection of the predator (e.g. Vásquez-Cruz, 2020), and parts of the prey not consumed and left by predators. (e.g. Silva *et al.*, 2018). Identification of the species is necessary for a better understanding of the dynamics in predator-prey relationships.

*Sceloporus bicanthalis* and *Abronia graminea* are lizard species endemic to Mexico. *Sceloporus bicanthalis* is a medium-sized terrestrial species (adult snout-vent length [SVL] ca. 60 mm) belonging to the family Phrynosomatidae (Canseco-Márquez and Gutiérrez-Mayén, 2010). It occurs in open areas of rocky grassland in forests or xeric scrub in the states of Hidalgo, Mexico, Oaxaca, Puebla, and Veracruz (Ramírez-Bautista *et al.*, 2014), and is considered common within its restricted distribution (Flores-Villela and Santos-Barrera, 2007a). *Abronia graminea* is a larger, mostly arboreal species (adult SVL ca. 110 mm) belonging to the family Anguidae. It is distributed in the highlands of central Veracruz,

southwestern Puebla, and northern Oaxaca, and inhabits a wide array of mesic forest types but often dominated by oak (*Quercus* spp.) (Canseco-Márquez and Gutiérrez-Mayén, 2010). The species is endangered by deforestation, climate change, and illegal harvest for pet trade, and is protected by Mexican law under the category Special Protection (IUCN Red List of Threatened Species, Flores-Villela and Santos-Barrera, 2007b; SEMARNAT 2010).

The Loggerhead Shrike (*Lanius ludovicianus*) is a passeriform bird in the family Laniidae with a wide distribution in North America, from southern Canada to northeastern Oaxaca in Mexico (Miller, 1931; Vázquez *et al.*, 2009). It has a carnivorous diet that includes invertebrates, amphibians, reptiles, rodents, and small birds (Jensen, 2013; Cogalniceanu, *et al.*, 2015). Its foraging behavior includes impaling its prey on thorns, branches, and even barbed wire, both to secure the prey for consumption and also for longer-term storage (Craig, 1978). Here we report for the first case of the predation on *Sceloporus bicanthalis* and *Abronia graminea* lizards presumably by the Loggerhead shrike (*Lanius ludovicianus*) in the central west of Veracruz, Mexico.

On 4 December 2019, we found seven lizards impaled on plant thorns, within a 50 m radius of each other, ca. 3 airline km north of Acultzingo

(18°44'30.7"N, 97°19'43.2"W, WGS 84, elev. 2520 m), Veracruz, Mexico. The area is characterised by pine-oak forest remnants, *Pinus patula* plantations, open areas of cattle pasture, and corn crops. We recorded two males and three females adults of *Sceloporus bicanthalis* and a skull of *Abronia graminea* impaled on thorns of unidentified shrubs at a height of 110–210 cm above the ground (Fig. 1A, B). We also found a male of *Sceloporus bicanthalis* on a terminal leaf spine of a maguey (*Agave* sp.) at a height of 150 cm above the ground (Fig. 1C).


Although we did not observe the predation events, we suggest that these *Sceloporus bicanthalis* and *Abronia graminea* were predated by *Lanius ludovicianus*, since it is a species observed in the study site (Fuentes-Moreno pers. comm., Howell and Webb, 1995) and whose foraging behavior includes the impaling of its prey (Yosef and Pinshowb, 2005). No other bird species in the study area exhibits this type of foraging behavior. Additionally, many species of lizards have been previously documented as preys of *L. ludovicianus* and were predated by impalement. Prey species include *Phrynosoma modestum* (Reid and Fulbright, 1981), *Anniella pulchra* (Fisher, 1901), *Anolis carolinensis*, *Plestiodon inexpectatus*, *Scincella lateralis*, *Aspidoscelis sexlineata* (Tyler, 1991; Yosef and Grubb, 1993), *Phrynosoma cornutum* (Tyler, 1991; Lemos-Espinal et al., 1998), *Sceloporus edbelli* (Lemos-Espinal et al., 2001), *Uma inornata* (Barrows, 2006), *Ophisaurus ventralis* (Clarke et al., 2011), *Dipsosaurus catalinensis* (Carbajal-Márquez et al., 2012), *Phrynosoma braconnieri* (Arias-Balderas et al., 2012), *Anniella grinnelli*, *Uta stansburiana*

(Herr and Papenfuss, 2017), *Phrynosoma hernandesi* (Cairns et al., 2017), *Anolis sagrei* (Simpson et al., 2019), *Phrynosoma mcallii* (e. g. Duncan, 1994; Lara-Resendiz et al., 2019), *Sceloporus aeneus* and *Plestiodon brevirostris* (Gómez-Campos et al., 2019). *Lanius ludovicianus* inhabits open areas where it usually hunts (Howell and Webb, 1995), and many of the lizard species known in its diet are terrestrial and inhabit open areas (e.g. species of Phrynosomatidae and Scincidae families), suggesting that feeding on *Sceloporus bicanthalis* may be recurrent, while predation on *Abronia graminea* could be a fortuitous case since it is an arboreal species (Canseco-Márquez and Gutiérrez-Mayen, 2010). Refuge availability for prey and abundance of both prey and predator species could influence these predator-prey interactions (e.g. Sih, 1987; Kumar, 2006; Das and Samanta, 2019).

Our observations are the first documented records of predation on *Abronia graminea* and add information about *Sceloporus bicanthalis* predators since only have been reported a snake species, *Crotalus intermedius*, as a predator (Schramer et al., 2020). These types of records are important for a better understanding of predator-prey dynamics, especially in reptiles species with secretive habits (e. g. Wiseman et al., 2018), threatened with extinction and restricted distribution, as *Abronia graminea*.

### Acknowledgments

We thank Alejandra Vásquez-Cruz for help in the field and for providing the photograph included in Figure 1C, and to Axel Fuentes-Moreno for the information about *L. ludovicianus*. We are also grateful



**Figure 1.** Impaled and predated lizards of *Sceloporus bicanthalis* and *Abronia graminea* in Acultzingo, Veracruz, Mexico. (A) *Abronia graminea* on thorn bush, (B) female *Sceloporus bicanthalis* on thorn bush, and (C) male *S. bicanthalis* on spine of *Agave* sp.

to two anonymous reviewers and Adam G. Clause for their valuable recommendations that improved the manuscript.

### Literature cited

- Arias-Balderas, S.F.; Hernandez-Rios, A. & Correa-Sanchez, F. 2012. *Phrynosoma braconneri* (Short-tailed Horned Lizard). Predation. *Herpetological Review* 43: 335–336.
- Barrows, C.W. 2006. Population dynamics of a threatened sand dune lizard. *The Southwestern Naturalist* 51: 514–523.
- Cairns, K.A.; Babineau, J. & Cairns, N.A. 2017. *Phrynosoma hernandesi* (Greater Short-Horned Lizard). Predation. *Herpetological review* 48: 853.
- Canseco-Márquez, L. & Gutiérrez-Mayén, M.G. 2010. Anfibios y reptiles del Valle de Tehuacán-Cuicatlán. Comisión Nacional para el conocimiento y uso de la biodiversidad, México D.F., México.
- Carbajal-Marquez, R.A.; Rodriguez-Canseco, J.M. & Arnaud, G. 2012. *Dipsosaurus catalinensis* (Santa Catalina Island Desert Iguana). Predation. *Herpetological Review* 43: 131–132.
- Clarke, F.O.; Jensen, J.B. & Crawford, J.A. 2011. *Ophisaurus ventralis* (Eastern glass lizard). Postmortem predation. *Herpetological Review* 42: 607.
- Cogalniceanu, D.; Valdeón, A.; Gosá, A.; Al-Hemaidi, A.A.M. & Castilla, M.A. 2015. Shrike predation on the lizard *Mesalina adramitana* in Qatar; a review of reported reptile and amphibian prey. *Qscience Connect* 2015: 1.
- Craig, R.B. 1978. An analysis of the predatory behavior of the Loggerhead Shrike. *Auk* 95: 221–234.
- Das, A., & Samanta, G.P. 2019. A prey-predator model with refuge for prey and additional food for predator in a fluctuating environment. *Physica A: Statistical Mechanics and Its Applications* 122844.
- Duncan, R.; Esque, T. & Echols, K. 1994. *Phrynosoma mcallii* (Flat-tailed Horned Lizard). Predation. *Herpetological Review* 25: 68.
- Fisher, W.K. 1901. Shrike notes. *Condor* 3: 48–49.
- Flores-Villela, O. & Santos-Barrera, G. 2007a. *Sceloporus bicanthalis*. The IUCN Red List of Threatened Species 2007: e.T64089A12736142. <https://dx.doi.org/10.2305/IUCN.UK.2007.RLTS.T64089A12736142.en>. Downloaded on 27 July 2020.
- Flores-Villela, O. & Santos-Barrera, G. 2007b. *Abronia graminea*. The IUCN Red List of Threatened Species 2007: e.T63678A12695490. <https://dx.doi.org/10.2305/IUCN.UK.2007.RLTS.T63678A12695490.en>. Downloaded on 27 July 2020.
- Gómez-Campos, J.E.; Domínguez-Godoy, M.A. & Díaz de la Vega-Pérez, A.H. 2019. Depredación de lagartijas por *Lanius ludovicianus* (Alcaudón americano) en el Parque Nacional la Malinche, México. *Revista Latinoamericana de Herpetología* 2: 85–87.
- Herr, M.W. & Papenfuss, T.J. 2017. *Anniella grinnelli* (Bakersfield Legless Lizard). Predation. *Herpetological Review* 48: 633.
- Howell, S.N.G. & Webb, S. 1995. *A guide to the birds of Mexico and Northern Central American*. New York (USA), Oxford University Press.
- Jensen, J.B. 2013. *Ambystoma tigrinum* (Tiger Salamander). Predation. *Herpetological Review* 34: 132–133.
- Kumar Kar, T. 2006. Modelling and analysis of a harvested prey-predator system incorporating a prey refuge. *Journal of Computational and Applied Mathematics* 185: 19–33.
- Lara-Resendiz, R.A.; Valdez-Villavicencio, J.H.; Pérez-Delgadillo, A.G.; Pinto-Santana, H.D. & Galina-Tessaro, P. 2019. Predation on Flat-tailed Horned Lizard (*Phrynosoma mcallii*) by Loggerhead Shrike (*Lanius ludovicianus*). *Revista Latinoamericana de Herpetología* 2: 44–47.
- Lemos-Espinal, J.A.; Chiszar, D.; Henke, C. & Smith, H.M. 1998. *Phrynosoma cornutum* (Texas Horned Lizard). Predation. *Herpetological Review* 29: 168.
- Lemos-Espinal, J.A.; Chiszar, D. & Smith, H.B. 2001. *Sceloporus belli* (Bell's Spiny Lizard). Predation. *Herpetological Review* 32: 42–43.
- Miller, A.H. 1931. *Systematic revision and natural history of the American Shrikes (Lanius)*. University of California Publications in Zoology.
- Ramírez-Bautista, A.; Hernández-Salinas, U.; Cruz-Elizalde, R.; Berriozabal-Islas, C.; Lara-Tufiño, D.; Mayer-Goyenechea I. G. & Castillo-Cerón, J. 2014. Los anfibios y reptiles de Hidalgo, México: Diversidad, Biogeografía y Conservación. Sociedad Herpetológica Mexicana, A. C.
- Reid, W.H. & Fulbright, H.J. 1981. Impaled prey of the Loggerhead Shrike in the Northern Chihuahuan Desert. *The Southwestern Naturalist* 26: 204–205.
- Schramer, T.D.; Kalki, Y.; West, T.R. & Wylie, D.B. 2020. Notes on the diet of *Crotalus intermedius* Troschel, 1865. *Revista Latinoamericana de Herpetología* 3: 11–115.
- Sih, A. 1987. Prey refuges and predator-prey stability. *Theoretical Population Biology* 31: 1–12.
- Silva Brito, E.; Miranda, E. & Tortato F.R. 2018. Chelonian predation by Jaguars (*Panthera onca*). *Chelonian Conservation and Biology* 17: 280–283.
- Simpson, S.E.; Giery, S.T.; Stroud, J.T. & Thawley, C.J. 2019. *Anolis sagrei* (Brown anole). Predation. *Herpetological Review* 50: 362–363.
- Steen, D.A.; McClure, C.J.W.; Sutton, W.B.; Rudolph, D.C., Pierce, J.B., Lee, J.R., Smith, L.L., Gregory, B.B., Baxley, D.L., Stevenson, D.J. & Guyer, C. 2014. Copperheads are common where kingsnakes are not: relationships between the abundances of a predator and one of their prey. *Herpetologica* 70: 69–76.
- Tyler, J.D. 1991. Vertebrate prey of the Loggerhead Shrike in Oklahoma. *Proceedings of the Oklahoma Academy of Science* 71: 17–20.
- Vásquez-Cruz, V. 2020. New prey records for the Atlantic Central American Milksnake *Lampropeltis polyzona* (Serpentes: Colubridae). *Phyllomedusa* 19: 107–111.
- Vásquez-Cruz, V. & Reynoso-Martínez, A. 2020. Contributions to the knowledge of the natural history of *Claudius angustatus* (Testudines: Kinosternidae) in Veracruz, Mexico. *Phyllomedusa* 19:113–116.
- Vázquez, L., Moya, H. & Arizmendi, M.D.C. 2009. Avifauna de la selva baja caducifolia en la cañada del río Sabino, Oaxaca, México. *Revista Mexicana de Biodiversidad* 80(2): 535–549.
- Wiseman, K.D.; Greene, H.W.; Koo, M.S. & Long, D.J. 2018. Feeding ecology of a generalist Predator, The California Kingsnake (*Lampropeltis californiae*): Why rare prey matter. *Herpetological Conservation and Biology* 14:1–30.
- Yosef, R. & Grubb, T.C. Jr. 1993. Effect of vegetation height

Vásquez-Cruz & Reynoso-Martínez — Predation of *Abronia graminea* and *Sceloporus bicanthalis*

on hunting behavior and diet of Loggerhead Shrikes. *The Condor* 95:127–31.

Yosef, R. & Pinshow, B. 2005. Impaling in true shrikes (Laniidae): A behavioral and ontogenic perspective. *Behavioural Processes* 69: 363–367.

© 2020 por los autores, licencia otorgada a la Asociación Herpetológica Argentina. Este artículo es de acceso abierto y distribuido bajo los términos y condiciones de una licencia Atribución-No Comercial 2.5 Argentina de Creative Commons. Para ver una copia de esta licencia, visite <http://creativecommons.org/licenses/by-nc/2.5/ar/>